

Medication Support Nurse Program

For patients prescribed Dysport and
enrolled in IPSEN CARES



Actor portrayal

Medication Support Nurses provide individualized patient support throughout the treatment journey.

Support offerings include:

- Multiple touchpoints during each injection cycle to address any patient questions
- Education about Dysport consistent with the US Prescribing Information
- Disease state education to help patients understand their condition
- Information to help patients understand how to work with their care team to set and achieve treatment goals
- Support to help patients receive their injections in a timely manner
- Continuity of care support to mitigate potential gaps in care

Medication Support Nurses are available Monday – Friday,
9:00 AM – 5:00 PM. Please call 1-866-435-5677.

SCAN THE QR CODE to enroll into IPSEN CARES and start receiving nursing support, or visit ipsencares.com/dysport



Indication and Important Safety Information about DYSPORT (abobotulinumtoxinA)

What is DYSPORT?

DYSPORT is a prescription medicine that is injected into muscles and used to treat:

- increased muscle stiffness in people 2 years of age and older with spasticity
- cervical dystonia in adults

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about DYSPORT?

DYSPORT may cause serious side effects that can be life threatening. These problems can happen within hours, or days to weeks after an injection of DYSPORT. Call your doctor or get medical help right away if you have any of these problems after treatment with DYSPORT:

Please see additional Important Safety Information on next page and accompanying full [Prescribing Information](#), including [Medication Guide](#) with IMPORTANT WARNING.

INFORMATION about DYSPORT® (abobotulinumtoxinA), continued

What is the most important information I should know about DYSPORT? (continued)

- **Problems swallowing, speaking, or breathing.** These problems can happen within hours to weeks after an injection of DYSPORT due to weakening of associated muscles. Death can happen as a complication if you have severe problems with swallowing or breathing after treatment with DYSPORT. Swallowing problems may last for several weeks. People with certain pre-existing swallowing or breathing problems have the highest risk of getting these problems.
- **Spread of toxin effects.** In some cases, the effect of botulinum toxin may affect areas of the body away from the injection site and cause symptoms of a serious condition called botulism including loss of strength and muscle weakness all over the body, double or blurred vision, drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, loss of bladder control, and trouble breathing or swallowing.

These symptoms can happen within hours to weeks after you receive an injection of DYSPORT. These problems could make it unsafe for you to drive a car, operate machinery, or do other dangerous activities.

Who should not receive DYSPORT?

Do not receive DYSPORT if you: are allergic to DYSPORT or any of its ingredients, or to cow's milk protein; had an allergic reaction to any other botulinum toxin product such as Myobloc®, Botox®, or Xeomin®; have a skin infection at the planned injection site.

What should I tell my doctor before receiving DYSPORT?

Tell your doctor about all your medical conditions, including if you:

- have a disease that affects your muscles and nerves, such as amyotrophic lateral sclerosis (ALS or Lou Gehrig's disease), myasthenia gravis, or Lambert-Eaton syndrome
- have allergies to any botulinum toxin product or had any side effect from any botulinum toxin product in the past
- have or have had problems with breathing (such as asthma or emphysema), swallowing, or bleeding
- have diabetes
- have or have had a slow heartbeat, or other problem with your heart rate or rhythm
- have plans to have surgery or had surgery on your face
- have weakness in or near your muscles being treated, drooping eyelids, experienced dry eye with previous use of botulinum toxin products, or any other change in the way your face normally looks
- are pregnant (or plan to be) or breast-feeding (or planning to)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal products. Using DYSPORT with certain other medicines may cause serious side effects.

Do not start any new medicines until you have told your doctor that you have received DYSPORT in the past.

Especially tell your doctor if you: have received any other botulinum toxin product in the last four months or any time in the past (be sure your doctor knows exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants, an allergy or cold medicine, or a sleep medicine.

What are the possible side effects of DYSPORT?

- **The most common side effects in adults with upper limb spasticity** include muscle weakness; **in adults with lower limb spasticity** include muscle weakness, pain in arms or legs, and falls
- **The most common side effects in children (2 to 17 years of age) with upper limb spasticity** include upper respiratory tract infection and sore throat; **in children with lower limb spasticity** include stuffy or runny nose, sore throat, cough, and fever
- **The most common side effects in people with cervical dystonia** include muscle weakness, muscle pain, difficulty swallowing, dry mouth, problems speaking, injection site pain or discomfort, feeling of tiredness, eye problems, and headache

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of DYSPORT. For more information, ask your doctor or pharmacist. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full [Prescribing Information](#), including [Medication Guide](#) with IMPORTANT WARNING.

DYSPORT is a registered trademark of Ipsen Biopharm Limited.

IPSEN CARES is a registered trademark of Ipsen S.A.

All other trademarks are property of their respective owners.

©2024 Ipsen Biopharmaceuticals, Inc. All Rights Reserved. October 2024 DYS-US-009134

